



Recommended Vaccination/Medication Protocol for High Exposure Kennels

- 2 Weeks: *Deworm with pyrantel pamoate (1cc per 5#)
- 4 Weeks: *Deworm with pyrantel pamoate (1cc per 5#)
*Give intranasal Bordetella bronchiseptica vaccination
- 4 ½ Weeks: *Vaccinate with Neopar (1cc under the skin)
- 6 Weeks: *Deworm with pyrantel pamoate (1cc per 5#)
*Vaccinate with Vanguard Plus 5
*Clean ears and trim nails. Treat for ear mites with Frontline Spray (1 pump per ear)
*+/- 7 days of Albon/Metro for coccidia/giardia prevention at weaning
- 7 Weeks: *Clean ears and treat for ear mites.
- 8 Weeks: *Deworm with Panacur for 3 days (1cc per 4#)
*Vaccinate with Vanguard Plus 5
*10 days of Albon/Metro for coccidia/giardia prevention
- 10 Weeks: *Deworm with pyrantel pamoate (1cc per 5#)
*Vaccinate with Vanguard Plus 5
- 12 Weeks: *Deworm with Panacur for 3 days (1cc per 4#)
*Vaccinate with Vanguard Plus 5
- 15 Weeks: *Deworm with pyrantel pamoate (1cc per 5#)
*Vaccinate with Vanguard Plus 5

*The following vaccination protocol is not a one size fits all protocol. Different situations may need a different protocol. Any **HIGH TITER** 5 way vaccination can be used in place of the Vanguard Plus 5. Please consult your veterinarian prior to changing your current protocol.